

**METHODS AND COMPOSITIONS USING  
THALIDOMIDE FOR THE TREATMENT  
AND MANAGEMENT OF CANCERS AND OTHER DISEASES**

**1. FIELD OF THE INVENTION**

This invention relates to methods of treating, preventing and/or managing specific  
5 cancers, and other diseases including, but not limited to, those associated with, or  
characterized by, undesired angiogenesis, by the administration of thalidomide alone or in  
combination with other therapeutics. In particular, the invention encompasses the use of  
specific combinations, or "cocktails," of drugs and other therapy, *e.g.*, radiation to treat  
these specific cancers, including those refractory to conventional therapy. The invention  
10 also relates to pharmaceutical compositions and dosing regimens.

**2. BACKGROUND OF THE INVENTION**

**2.1 PATHOBIOLOGY OF CANCER AND OTHER DISEASES**

Cancer is characterized primarily by an increase in the number of abnormal cells  
derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, or  
15 lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant  
sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a  
multistep process that begins with minor preneoplastic changes, which may under certain  
conditions progress to neoplasia. The neoplastic lesion may evolve clonally and develop an  
increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under  
20 conditions in which the neoplastic cells escape the host's immune surveillance. Roitt, I.,  
Brostoff, J and Kale, D., *Immunology*, 17.1-17.12 (3rd ed., Mosby, St. Louis, Mo., 1993).

There is an enormous variety of cancers which are described in detail in the medical  
literature. Examples includes cancer of the lung, colon, rectum, prostate, breast, brain, and  
intestine. The incidence of cancer continues to climb as the general population ages, as new  
25 cancers develop, and as susceptible populations (*e.g.*, people infected with AIDS or  
excessively exposed to sunlight) grow. A tremendous demand therefore exists for new  
methods and compositions that can be used to treat patients with cancer.

Many types of cancers are associated with new blood vessel formation, a process  
known as angiogenesis. Several of the mechanisms involved in tumor-induced  
30 angiogenesis have been elucidated. The most direct of these mechanisms is the secretion by  
the tumor cells of cytokines with angiogenic properties. Examples of these cytokines

include acidic and basic fibroblastic growth factor (a,b-FGF), angiogenin, vascular endothelial growth factor (VEGF), and TNF- $\alpha$ . Alternatively, tumor cells can release angiogenic peptides through the production of proteases and the subsequent breakdown of the extracellular matrix where some cytokines are stored (e.g., b-FGF). Angiogenesis can also be induced indirectly through the recruitment of inflammatory cells (particularly macrophages) and their subsequent release of angiogenic cytokines (e.g., TNF- $\alpha$ , bFGF).

A variety of other diseases and disorders are also associated with, or characterized by, undesired angiogenesis. For example, enhanced or unregulated angiogenesis has been implicated in a number of diseases and medical conditions including, but not limited to, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, rubeosis (neovascularization of the angle), viral diseases, genetic diseases, inflammatory diseases, allergic diseases, and autoimmune diseases. Examples of such diseases and conditions include, but are not limited to: diabetic retinopathy; retinopathy of prematurity; corneal graft rejection; neovascular glaucoma; retrolental fibroplasia; and proliferative vitreoretinopathy.

Accordingly, compounds that can control angiogenesis or inhibit the production of certain cytokines, including TNF- $\alpha$ , may be useful in the treatment and prevention of various diseases and conditions.

## **2.2 METHODS OF TREATING CANCER**

Current cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (see, for example, Stockdale, 1998, *Medicine*, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of a patient or may be unacceptable to the patient. Additionally, surgery may not completely remove neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue. Radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent. Although hormonal therapy can be effective, it is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of cancer cells. Biological therapies and immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division. Gilman et al., *Goodman and Gilman's: The Pharmacological Basis of Therapeutics*, Tenth Ed. (McGraw Hill, New York).


Despite availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Stockdale, *Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10, 1998. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous side effects including severe nausea, bone marrow depression, and immunosuppression. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even if those agents act by different mechanism from those of the drugs used in the specific treatment. This phenomenon is referred to as pleiotropic drug or multidrug resistance. Because of the drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Other diseases or conditions associated with, or characterized by, undesired angiogenesis are also difficult to treat. However, some compounds such as protamine, heparin and steroids have been proposed to be useful in the treatment of certain specific diseases. Taylor et al., *Nature* 297:307 (1982); Folkman et al., *Science* 221:719 (1983); and U.S. Pat. Nos. 5,001,116 and 4,994,443. Thalidomide and certain derivatives of it have also been proposed for the treatment of such diseases and conditions. U.S. patent nos. 5,593,990, 5,629,327, 5,712,291, 6,071,948 and 6,114,355 to D'Amato.

Still, there is a significant need for safe and effective methods of treating, preventing and managing cancer and other diseases and conditions, particularly for diseases that are refractory to standard treatments, such as surgery, radiation therapy, chemotherapy and hormonal therapy, while reducing or avoiding the toxicities and/or side effects associated with the conventional therapies.

### **2.3 THALIDOMIDE**

Thalidomide is a racemic compound sold under the trade name Thalomid® and chemically named  $\alpha$ -(N-phthalimido)glutarimide or 2-(2,6-dioxo-3-piperidiny)-1H-isindole-1,3(2H)-dione. The compound has structure I:



I

Thalidomide was originally developed in the 1950's to treat morning sickness, but due to its teratogenic effects was withdrawn from use. Thalidomide has been approved in the United States for the acute treatment of the cutaneous manifestations of erythema nodosum leprosum in leprosy. *Physicians' Desk Reference*, 1153-1157 (57th ed., 2003). Because its administration to pregnant women can cause birth defects, the sale of thalidomide is strictly controlled. *Id.* Thalidomide has reportedly been studied in the treatment of other diseases, such as chronic graft-vs-host disease, rheumatoid arthritis, sarcoidosis, several inflammatory skin diseases, and inflammatory bowel disease. *See generally*, Koch, H.P., *Prog. Med. Chem.* 22:165-242 (1985). *See also*, Moller, D.R., *et al.*, *J. Immunol.* 159:5157-5161 (1997); Vasiliauskas, E.A., *et al.*, *Gastroenterology* 117:1278-1287 (1999); Ehrenpreis, E.D., *et al.*, *Gastroenterology* 117:1271-1277 (1999). It has further been alleged that thalidomide can be combined with other drugs to treat ischemia/reperfusion associated with coronary and cerebral occlusion. *See* U.S. Patent No. 5,643,915, which is incorporated herein by reference.

Thalidomide has reportedly been clinically investigated in the treatment of specific types of cancers, such as refractory multiple myeloma, brain, melanoma, breast, colon, mesothelioma, and renal cell carcinoma. *See, e.g.*, Singhal, S., *et al.*, *New England J. Med.* 341(21):1565-1571 (1999); and Marx, G.M., *et al.*, *Proc. Am. Soc. Clin. Oncology* 18:454a (1999). It has further been reported that thalidomide can be used to prevent the development of chronic cardiomyopathy in rats caused by doxorubicin. Costa, P.T., *et al.*, *Blood* 92(10:suppl. 1):235b (1998). Other reports concerning the use of thalidomide in the treatment of specific cancers include its combination with carboplatin in the treatment of glioblastoma multiforme. McCann, J., *Drug Topics* 41-42 (June 21, 1999). Thalidomide has also reportedly been used as an antiemetic during the treatment of astrocytoma. Zwart, D., *Arzneim.-Forsch.* 16(12):1688-1689 (1966).

If there is a general mechanism by which thalidomide aids in the treatment of some cancers, its nature remains unclear. *See, e.g.*, Moreira, A.L., *et al.*, *J. Exper. Med.* 177:1675-1680 (1993); McHugh, S.M., *et al.*, *Clin. Exper. Immunol.* 99:160-167 (1995); and Moller, D.R., *et al.*, *J. Immunol.* 159:5157-5161 (1997). It has been reported, however, that

thalidomide is an antiangiogenic agent that can suppress tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) and interleukin 12 (IL-12) production. *See, e.g., Moller, D.R., et al., J. Immunol.* 159:5157-5161 (1997); Moreira, A.L., *et al., J. Exp. Med.* 177:1675-1680 (1993); U.S. Patent Nos. 5,593,990, 5,629,327, and 5,712,291 to D'Amato and U.S. Patent No. 5,385,901 to Kaplan.

- 5 And *in vitro* studies suggest that thalidomide affects the production of a variety of other proteins. *See, e.g., McHugh, S.M., et al., Clin. Exp. Immunol.* 99:160-167 (1995).  
Thalidomide may also affect mechanisms related to epithelial or endothelial function or growth. D'Amato M., *et al., Proc. Natl. Acad. Sci.* 91:4082-4085(1994).

### 3. SUMMARY OF THE INVENTION

- 10 This invention encompasses methods of treating and preventing certain types of cancer, including primary and metastatic cancer, as well as cancers that are refractory or resistant to conventional chemotherapy. The methods comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate,  
15 stereoisomer, clathrate, or prodrug thereof. The invention also encompasses methods of managing certain cancers (*e.g.*, preventing or prolonging their recurrence, or lengthening the time of remission) which comprise administering to a patient in need of such management a prophylactically effective amount of thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

- 20 In particular methods of the invention, thalidomide is administered in combination with a therapy conventionally used to treat, prevent or manage cancer. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

- This invention also encompasses methods of treating, managing or preventing  
25 diseases and disorders other than cancer that are associated with, or characterized by, undesired angiogenesis, which comprise administering to a patient in need of such treatment, management or prevention a therapeutically or prophylactically effective amount of thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

- 30 In other methods of the invention, thalidomide is administered in combination with a therapy conventionally used to treat, prevent or manage diseases or disorders associated with, or characterized by, undesired angiogenesis. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention encompasses pharmaceutical compositions, single unit dosage forms, dosing regimens and kits which comprise thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second, or additional, active agent. Second active agents include specific combinations, or "cocktails,"  
5 of drugs.

#### 4. DETAILED DESCRIPTION OF THE INVENTION

A first embodiment of the invention encompasses methods of treating, managing, or preventing cancer which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of thalidomide, or a  
10 pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods encompassed by this embodiment, thalidomide is administered in combination with another drug ("second active agent") or method of treating, managing, or preventing cancer. Second active agents include small molecules and large molecules  
15 (*e.g.*, proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of thalidomide include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage cancer.

20 Another embodiment of the invention encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are characterized by undesired angiogenesis. These methods comprise the administration of a therapeutically or prophylactically effective amount of thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

25 Examples of diseases and disorders associated with, or characterized by, undesired angiogenesis include, but are not limited to, inflammatory diseases, autoimmune diseases, viral diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, and rubeosis (neovascularization of the angle). Specific examples of the diseases and disorders  
30 associated with, or characterized by, undesired angiogenesis include, but are not limited to, endotoxemia, toxic shock syndrome, osteoarthritis, retrovirus replication, wasting, meningitis, silica-induced fibrosis, asbestos-induced fibrosis, veterinary disorder, malignancy-associated hypercalcemia, stroke, circulatory shock, periodontitis, gingivitis, macrocytic anemia, refractory anemia, and 5q-syndrome.

In particular methods encompassed by this embodiment, thalidomide is administered in combination with a second active agent or method of treating, managing, or preventing the disease or condition. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells.

5 Methods, or therapies, that can be used in combination with the administration of thalidomide include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage disease and conditions associated with, or characterized by, undesired angiogenesis.

10 The invention also encompasses pharmaceutical compositions (e.g., single unit dosage forms) that can be used in methods disclosed herein. Particular pharmaceutical compositions comprise thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second active agent.

#### 4.1 COMPOUNDS OF THE INVENTION

15 Compounds used in the invention include racemic thalidomide, stereomerically enriched or stereomerically pure thalidomide, and pharmaceutically acceptable salts, solvates, hydrates, stereoisomers, clathrates, and prodrugs thereof.

As used herein and unless otherwise indicated, the term "stereomerically pure" means a composition that comprises one stereoisomer of a compound and is substantially  
20 free of other stereoisomer of that compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomer of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomer of the compound, even more preferably greater than about  
25 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomer of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomer of the compound.

As used herein and unless otherwise indicated, the term "stereomerically enriched"  
30 means a composition that comprises greater than about 60% by weight of one stereoisomer of a compound, preferably greater than about 70% by weight, more preferably greater than about 80% by weight of one stereoisomer of the compound.

As used herein and unless otherwise indicated, the term "enantiomerically pure" means a stereomerically pure composition of a compound having one chiral center.

Similarly, the term "enantiomerically enriched" means a stereomerically enriched composition of a compound having one chiral center.

Thalidomide can either be commercially purchased (from Celgene Corp., New Jersey) or prepared according to the known methods. *See, e.g., I. D. Fratta et al., Toxicol. Appl. Pharmacol.* 7, 268 (1965), and the references disclosed therein. Enantiomerically pure thalidomide can be resolved using known resolving agents or chiral columns as well as other standard synthetic organic chemistry techniques. *See, e.g., Blaschke, Arzneimittelforschung* 29: 1640-1642 (1979); Shealy *et al., Chem. Indus.* 1030 (1965); and Casini *et al., Farmaco Ed. Sci.* 19:563 (1964).

As used herein, unless otherwise specified, the term "pharmaceutically acceptable salt(s)" includes salts of acidic or basic moieties of the compound(s) to which the term refers. Basic moieties are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that can be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, *i.e.,* salts containing pharmacologically acceptable anions. Suitable organic acids include, but are not limited to, maleic, fumaric, benzoic, ascorbic, succinic, acetic, formic, oxalic, propionic, tartaric, salicylic, citric, gluconic, lactic, mandelic, cinnamic, oleic, tannic, aspartic, stearic, palmitic, glycolic, glutamic, gluconic, glucaronic, saccharic, isonicotinic, methanesulfonic, ethanesulfonic, p-toluenesulfonic, benzenesulfonic acids, or pamoic (*i.e.,* 1,1'-methylene-bis-(2-hydroxy-3-naphthoate) acids. Suitable inorganic acids include, but are not limited to, hydrochloric, hydrobromic, hydroiodic, sulfuric, phosphoric, or nitric acids. Compounds that include an amine moiety can form pharmaceutically acceptable salts with various amino acids, in addition to the acids mentioned above.

Compounds that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds are those that form non-toxic base addition salts, *i.e.,* salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.

As used herein to describe a compound or chemical moiety, the term "derivative" means a compound or chemical moiety wherein the degree of saturation of at least one bond has been changed (*e.g.,* a single bond has been changed to a double or triple bond) or



wherein at least one hydrogen atom is replaced with a different atom or a chemical moiety. Examples of different atoms and chemical moieties include, but are not limited to, halogen, oxygen, nitrogen, sulfur, hydroxy, methoxy, alkyl, amine, amide, ketone, and aldehyde.

As used herein and unless otherwise indicated, the term "prodrug" means a  
5 derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (*in vitro* or *in vivo*) to provide the compound. Examples of prodrugs include, but are not limited to, derivatives of thalidomide that include biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate  
10 analogues. Other examples of prodrugs include derivatives of thalidomide that include -NO, -NO<sub>2</sub>, -ONO, or -ONO<sub>2</sub> moieties.

As used herein and unless otherwise indicated, the terms "biohydrolyzable carbamate," "biohydrolyzable carbonate," "biohydrolyzable ureide," "biohydrolyzable phosphate" mean a carbamate, carbonate, ureide, or phosphate, respectively, of a compound  
15 that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties *in vivo*, such as uptake, duration of action, or onset of action; or 2) is biologically inactive but is converted *in vivo* to the biologically active compound. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, aminoacids, hydroxyalkylamines,  
20 heterocyclic and heteroaromatic amines, and polyether amines.

As used herein and unless otherwise indicated, the term "biohydrolyzable ester" means an ester of a compound that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties *in vivo*, such as uptake, duration of action, or onset of action; or 2) is biologically inactive but is  
25 converted *in vivo* to the biologically active compound. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, alkoxyacyloxy esters, alkyl acylamino alkyl esters, and choline esters.

As used herein and unless otherwise indicated, the term "biohydrolyzable amide" means an amide of a compound that either: 1) does not interfere with the biological activity  
30 of the compound but can confer upon that compound advantageous properties *in vivo*, such as uptake, duration of action, or onset of action; or 2) is biologically inactive but is converted *in vivo* to the biologically active compound. Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides,  $\alpha$ -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides.

## 4.2 SECOND ACTIVE AGENTS

Thalidomide can be combined with other pharmacologically active compounds ("second active agents") in methods and compositions of the invention. It is believed that certain combinations work synergistically in the treatment of particular types of cancer and certain diseases and conditions associated with, or characterized by, undesired angiogenesis. Thalidomide can also work to alleviate adverse effects associated with certain second active agents, and some second active agents can be used to alleviate adverse effects associated with thalidomide.

One or more second active ingredients or agents can be used in the methods and compositions of the invention together with thalidomide. Second active agents can be large molecules (*e.g.*, proteins) or small molecules (*e.g.*, synthetic inorganic, organometallic, or organic molecules).

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Specific examples of the active agents are anti-CD40 monoclonal antibodies (such as, for example, SGN-40); histone deacetylase inhibitors (such as, for example, SAHA and LAQ 824); heat-shock protein-90 inhibitors (such as, for example, 17-AAG); insulin-like growth factor-1 receptor kinase inhibitors; vascular endothelial growth factor receptor kinase inhibitors (such as, for example, PTK787); insulin growth factor receptor inhibitors; lysophosphatidic acid acyltransferase inhibitors; I $\kappa$ B kinase inhibitors; p38MAPK inhibitors; EGFR inhibitors (such as, for example, gefitinib and erlotinib HCL); HER-2 antibodies (such as, for example, trastuzumab (Herceptin<sup>®</sup>) and pertuzumab (Omnitarg<sup>™</sup>)); VEGFR antibodies (such as, for example, bevacizumab (Avastin<sup>™</sup>)); VEGFR inhibitors (such as, for example, flk-1 specific kinase inhibitors, SU5416 and ptk787/zk222584); P13K inhibitors (such as, for example, wortmannin); C-Met inhibitors (such as, for example, PHA-665752); monoclonal antibodies (such as, for example, rituximab (Rituxan<sup>®</sup>), tositumomab (Bexxar<sup>®</sup>), edrecolomab (Panorex<sup>®</sup>) and G250); and anti-TNF- $\alpha$  antibodies.

Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells *in vitro* or *in vivo*. Others stimulate the division and differentiation of committed erythroid progenitors in cells *in vitro* or *in vivo*. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including

recombinant IL-II ("rIL2") and canarypox IL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alfa-n1, interferon alfa-n3, interferon beta-I a, and interferon gamma-I b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions of the invention include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, CA); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, WA); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, CA).

Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. patent nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. patent nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

This invention encompasses the use of native, naturally occurring, and recombinant proteins. The invention further encompasses mutants and derivatives (*e.g.*, modified forms) of naturally occurring proteins that exhibit, *in vivo*, at least some of the pharmacological activity of the proteins upon which they are based. Examples of mutants include, but are not limited to, proteins that have one or more amino acid residues that differ from the corresponding residues in the naturally occurring forms of the proteins. Also encompassed by the term "mutants" are proteins that lack carbohydrate moieties normally present in their naturally occurring forms (*e.g.*, nonglycosylated forms). Examples of derivatives include, but are not limited to, pegylated derivatives and fusion proteins, such as proteins formed by fusing IgG1 or IgG3 to the protein or active portion of the protein of interest. *See, e.g.*, Penichet, M.L. and Morrison, S.L., *J. Immunol. Methods* 248:91-101 (2001).

Large molecule active agents may be administered in the form of anti-cancer vaccines. For example, vaccines that secrete, or cause the secretion of, cytokines such as IL-2, G-CSF, and GM-CSF can be used in the methods, pharmaceutical compositions, and kits of the invention. *See, e.g.*, Emens, L.A., *et al.*, *Curr. Opinion Mol. Ther.* 3(1):77-84 (2001).

In one embodiment of the invention, the large molecule active agent reduces, eliminates, or prevents an adverse effect associated with the administration of thalidomide. Depending on the disease or disorder being treated, adverse effects can include, but are not limited to, drowsiness and somnolence, dizziness and orthostatic hypotension, neutropenia,

infections that result from neutropenia, increased HIV-viral load, bradycardia, Stevens-Johnson Syndrome and toxic epidermal necrolysis, and seizures (*e.g.*, grand mal convulsions). A specific adverse effect is neutropenia.

Second active agents that are small molecules can also be used to alleviate adverse effects associated with the administration of thalidomide. However, like some large molecules, many are believed to be capable of providing a synergistic effect when administered with (*e.g.*, before, after or simultaneously) thalidomide. Examples of small molecule second active agents include, but are not limited to, anti-cancer agents, antibiotics, immunosuppressive agents, and steroids.

Examples of anti-cancer agents include, but are not limited to: semaxanib; cyclosporin; etanercept; doxycycline; bortezomib; acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; broprimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; celecoxib (COX-2 inhibitor); chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epiropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iproplatin; irinotecan; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedopa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine;

peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprime; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin;

5 spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricirbine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard;

10 uredepa; vaporeotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; and zorubicin hydrochloride.

Other anti-cancer drugs include, but are not limited to: 20-epi-1,25

15 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen;

20 antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauosporine; beta lactam derivatives; beta-alethine; betaclamycin

25 B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B;

30 cetorelix; chlorlins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatin; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrodidemnin

B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziqune;  
 didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-;  
 dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine;  
 doxorubicin; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine;  
 5 edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine  
 analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate;  
 exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol;  
 flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex;  
 formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine;  
 10 ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin;  
 hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene;  
 idramantone; ilmofofosine; ilomastat; imatinib (*e.g.*, Gleevec<sup>®</sup>), imiquimod; immunostimulant  
 peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons;  
 interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine;  
 15 isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N  
 triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole;  
 leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone;  
 leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide  
 peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine;  
 20 lometrexol; lonidamine; losoxantrone; loxoribine; lurtotecan; lutetium texaphyrin;  
 lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin;  
 matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin;  
 methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim;  
 mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth  
 25 factor-saporin; mitoxantrone; mofarotene; molgramostim; Erbitux, human chorionic  
 gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; mustard  
 anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone;  
 N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine;  
 napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; nilutamide;  
 30 nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; oblimersen  
 (Genasense<sup>®</sup>); O<sup>6</sup>-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone;  
 ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin;  
 oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine;  
 palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine;

- pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds;
- 5 platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists;
- 10 raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; sizofiran; sobuzoxane;
- 15 sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors;
- 20 temoporfin; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors;
- 25 tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer.

- Other second active agents include, but are not limited to, mTOR inhibitors (such as
- 30 rapamycin), androgens, pamidronate, D1/3-MAGE3 peptide vaccine, NY-ESO-1/ISCOMATRIX vaccine, proteasome inhibitor PS-341, etanercept, mesna, bortezomib, azacytidine, and glucocorticoids.

Specific second active agents include, but are not limited to, 2-methoxyestradiol, telomestatin, inducers of apoptosis in multiple myeloma cells (such as, for example,

TRAIL), statins, semaxanib, cyclosporin, etanercept, doxycycline, bortezomib, oblimersen (Genasense<sup>®</sup>), remicade, docetaxel, celecoxib, melphalan, dexamethasone (Decadron<sup>®</sup>), steroids, gemcitabine, cisplatin, temozolomide, etoposide, cyclophosphamide, temodar, carboplatin, procarbazine, gliadel, tamoxifen, topotecan, methotrexate, Arisa<sup>®</sup>, taxol, taxotere, fluorouracil, leucovorin, irinotecan, xeloda, CPT-11, interferon alpha, pegylated interferon alpha (*e.g.*, PEG INTRON-A), capecitabine, cisplatin, thiotepa, fludarabine, carboplatin, liposomal daunorubicin, cytarabine, doxorubicin, paclitaxel, vinblastine, IL-2, GM-CSF, dacarbazine, vinorelbine, zoledronic acid, palmitronate, biaxin, busulphan, prednisone, bisphosphonate, arsenic trioxide, vincristine, doxorubicin (Doxil<sup>®</sup>), paclitaxel, ganciclovir, adriamycin, estramustine sodium phosphate (Emcyt<sup>®</sup>), sulindac, and etoposide.

### **4.3 METHODS OF TREATMENTS AND PREVENTION**

Methods of this invention encompass methods of treating, preventing and/or managing various types of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis. As used herein, unless otherwise specified, the term “treating” refers to the administration of a compound of the invention or other additional active agent after the onset of symptoms of the particular disease or disorder. As used herein, unless otherwise specified, the term “preventing” refers to the administration prior to the onset of symptoms, particularly to patients at risk of cancer, and other diseases and disorders associated with, or characterized by, undesired angiogenesis. The term “prevention” includes the inhibition of a symptom of the particular disease or disorder. Patients with familial history of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis are preferred candidates for preventive regimens. As used herein and unless otherwise indicated, the term “managing” encompasses preventing the recurrence of the particular disease or disorder in a patient who had suffered from it, and/or lengthening the time a patient who had suffered from the disease or disorder remains in remission.

As used herein, the term “cancer” includes, but is not limited to, solid tumors and blood born tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, spinal cord tumors, pheochromocytoma, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor



- prognosis malignant brain tumor, malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi's sarcoma, karotype acute myeloblastic leukemia,
- 5 Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, localized or metastatic melanoma (of any kind, including, but not limited to, ocular), peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma,
- 10 fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unrescectable hepatocellular carcinoma, Waldenstrom's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer,
- 15 papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation.

- As used herein to refer to diseases and conditions other than cancer, the terms "diseases or disorders associated with, or characterized by, undesired angiogenesis,"
- 20 "diseases or disorders associated with undesired angiogenesis," and "diseases or disorders characterized by undesired angiogenesis" refer to diseases, disorders and conditions that are caused, mediated or attended by undesired, unwanted or uncontrolled angiogenesis, including, but not limited to, inflammatory diseases, autoimmune diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular
- 25 diseases, and retina neovascular diseases.

- Examples of such diseases or disorders associated with undesired angiogenesis include, but are not limited to, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, proliferative vitreoretinopathy, trachoma, myopia, optic pits, epidemic keratoconjunctivitis, atopic keratitis, superior limbic
- 30 keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, lipid degeneration, bacterial ulcer, fungal ulcer, Herpes simplex infection, Herpes zoster infection, protozoan infection, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, mariginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, scleritis, Steven's Johnson disease, periphigoid radial

keratotomy, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis, chronic vitritis, Lyme's disease, Eales disease, Bechet's disease, retinitis, choroiditis, presumed ocular histoplasmosis, Bests disease, Stargarts disease, pars planitis, chronic retinal detachment, 5 hyperviscosity syndromes, toxoplasmosis, rubeosis, sarcodiosis, sclerosis, soriatis, psoriasis, primary sclerosing cholangitis, proctitis, primary biliary srosis, idiopathic pulmonary fibrosis, alcoholic hepatitis, endotoxemia, toxic shock syndrome, osteoarthritis, retrovirus replication, wasting, meningitis, silica-induced fibrosis, asbestos-induced fibrosis, malignancy-associated hypercalcemia, stroke, circulatory shock, periodontitis, gingivitis, 10 macrocytic anemia, refractory anemia, 5q- syndrome, and veterinary disorder caused by feline immunodeficiency virus, equine infectious anemia virus, caprine arthritis virus, visna virus, maedi virus or lenti virus.

In specific embodiments of the invention, diseases or disorders associated with undesired angiogenesis do not include congestive heart failure, cardiomyopathy, pulmonary 15 edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, HIV, hepatitis, adult respiratory distress syndrome, bone-resorption disease, chronic obstructive pulmonary diseases, chronic pulmonary inflammatory disease, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, fibrotic disease, cachexia, graft rejection, 20 rheumatoid spondylitis, osteoporosis, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythrematosus, erythema nodosum leprosum in leprosy, radiation damage, asthma, hyperoxic alveolar injury, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

This invention encompasses methods of treating patients who have been previously 25 treated for cancer or diseases or disorders associated with, or characterized by, undesired angiogenesis, but are non-responsive to standard therapies, as well as those who have not previously been treated. The invention also encompasses methods of treating patients regardless of patient's age, although some diseases or disorders are more common in certain age groups. The invention further encompasses methods of treating patients who have 30 undergone surgery in an attempt to treat the disease or condition at issue, as well as those who have not. Because patients with cancer and diseases and disorders characterized by undesired angiogenesis have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific

secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with cancer and other diseases or disorders.

Methods encompassed by this invention comprise administering one or more thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, to a patient (*e.g.*, a human) suffering, or likely to suffer, from cancer or a disease or disorder mediated by undesired angiogenesis.

In one embodiment of the invention, thalidomide is administered orally and daily in an amount of from about 50 to about 2,000 mg, preferably from about 50 to about 1,000 mg, and more preferably from about 50 to 800 mg. In a preferred embodiment, the recommended dose of thalidomide is from about 200 mg to about 800 mg.

In a specific embodiment, thalidomide is administered in an amount of from about 200 mg to about 800 mg per day to patients with relapsed multiple myeloma. In another specific embodiment, thalidomide is administered in an amount of from about 200 mg to about 800 mg to patients with solid tumor. In a particular embodiment, thalidomide is administered to patients with glioma.

In a specific embodiment, thalidomide is administered in an amount of from about 200 mg to about 800 mg per day, or alternatively every other day, to patients with diseases or disorders associated with, or characterized by, undesired angiogenesis including, but not limited to, endotoxemia, toxic shock syndrome, osteoarthritis, retrovirus replication, wasting, meningitis, silica-induced fibrosis, asbestos-induced fibrosis, veterinary disorder, malignancy-associated hypercalcemia, stroke, circulatory shock, periodontitis, gingivitis, macrocytic anemia, refractory anemia, and 5q- syndrome.

#### **4.3.1 COMBINATION THERAPY**

Specific methods of the invention comprise administering thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in combination with one or more second active agents, and/or in combination with radiation therapy, blood transfusions, or surgery. Examples of second active agents are disclosed herein (*see, e.g.*, section 4.2).

Administration of thalidomide and the second active agents to a patient can occur simultaneously or sequentially by the same or different routes of administration. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (*e.g.*, whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. A preferred route of administration for thalidomide is oral. Preferred routes of administration for the

second active agents or ingredients of the invention are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56<sup>th</sup> ed., 2002).

In one embodiment of the invention, the second active agent is administered intravenously or subcutaneously and once or twice daily in an amount of from about 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. The specific amount of the second active agent will depend on the specific agent used, the type of disease being treated or managed, the severity and stage of disease, and the amount(s) of thalidomide and any optional additional active agents concurrently administered to the patient. In a particular embodiment, the second active agent is oblimersen (Genasense<sup>®</sup>), GM-CSF, G-CSF, EPO, taxotere, irinotecan, dacarbazine, transretinoic acid, topotecan, pentoxifylline, ciprofloxacin, dexamethasone, vincristine, doxorubicin, COX-2 inhibitor, IL2, IL8, IL18, IFN, Ara-C, vinorelbine, or a combination thereof.

In certain embodiments, the second active agent is not a topoisomerase inhibitor (e.g., irinotecan). In certain embodiments, the second active agent is not temozolomide.

In a particular embodiment, GM-CSF, G-CSF or EPO is administered subcutaneously during about five days in a four or six week cycle in an amount of from about 1 to about 750 mg/m<sup>2</sup>/day, preferably in an amount of from about 25 to about 500 mg/m<sup>2</sup>/day, more preferably in an amount of from about 50 to about 250 mg/m<sup>2</sup>/day, and most preferably in an amount of from about 50 to about 200 mg/m<sup>2</sup>/day. In a certain embodiment, GM-CSF may be administered in an amount of from about 60 to about 500 mcg/m<sup>2</sup> intravenously over 2 hours, or from about 5 to about 12 mcg/m<sup>2</sup>/day subcutaneously. In a specific embodiment, G-CSF may be administered subcutaneously in an amount of about 1 mcg/kg/day initially and can be adjusted depending on rise of total granulocyte counts. The maintenance dose of G-CSF may be administered in an amount of about 300 (in smaller patients) or 480 mcg subcutaneously. In a certain embodiment, EPO may be administered subcutaneously in an amount of 10,000 units three times per week.

In another embodiment, thalidomide in an amount of from about 200 mg to about 800 mg per day and dacarbazine in an amount of about from 200 to 1,000 mg/m<sup>2</sup>/d are administered to patients with metastatic melanoma. In a specific embodiment, thalidomide is administered to patients with metastatic melanoma whose disease has progressed on treatment with dacarbazine, IL-2 or IFN. In a specific embodiment, thalidomide is administered to patients with relapsed or refractory multiple myeloma in combination with dexamethasone.

In another embodiment, thalidomide is administered with melphalan and dexamethasone to patients with amyloidosis. In a specific embodiment, thalidomide and steroids can be administered to patients with amyloidosis.

In another embodiment, thalidomide is administered with gemcitabine and  
5 cisplatin to patients with locally advanced or metastatic transitional cell bladder cancer.

In another embodiment, thalidomide is administered in combination with a second active ingredient as follows: temozolomide to pediatric patients with relapsed or progressive brain tumors or recurrent neuroblastoma; celecoxib, etoposide and cyclophosphamide for relapsed or progressive CNS cancer; temodar to patients with  
10 recurrent or progressive meningioma, malignant meningioma, hemangiopericytoma, multiple brain metastases, relapsed brain tumors, or newly diagnosed glioblastoma multiforms; irinotecan to patients with recurrent glioblastoma; carboplatin to pediatric patients with brain stem glioma; procarbazine to pediatric patients with progressive malignant gliomas; cyclophosphamide to patients with poor prognosis malignant brain  
15 tumors, newly diagnosed or recurrent glioblastoma multiforms; Gliadel® for high grade recurrent malignant gliomas; temozolomide and tamoxifen for anaplastic astrocytoma; or topotecan for gliomas, glioblastoma, anaplastic astrocytoma or anaplastic oligodendroglioma.

In another embodiment, thalidomide is administered with methotrexate and  
20 cyclophosphamide to patients with metastatic breast cancer.

In another embodiment, thalidomide is administered with temozolomide to patients with neuroendocrine tumors.

In another embodiment, thalidomide is administered with gemcitabine to patients with recurrent or metastatic head or neck cancer. In another embodiment, thalidomide is  
25 administered with gemcitabine to patients with pancreatic cancer.

In another embodiment, thalidomide is administered to patients with colon cancer in combination with Arisa®, taxol and/or taxotere.

In another embodiment, thalidomide is administered with capecitabine to patients with refractory colorectal cancer or patients who fail first line therapy or have poor  
30 performance in colon or rectal adenocarcinoma.

In another embodiment, thalidomide is administered in combination with fluorouracil, leucovorin, and irinotecan to patients with Dukes C & D colorectal cancer or to patients who have been previously treated for metastatic colorectal cancer.

In another embodiment, thalidomide is administered to patients with refractory colorectal cancer in combination with capecitabine, xeloda, and/or CPT-11.

5 In another embodiment, thalidomide is administered with capecitabine and irinotecan to patients with refractory colorectal cancer or to patients with unresectable or metastatic colorectal carcinoma.

In another embodiment, thalidomide is administered alone or in combination with interferon alpha or capecitabine to patients with unresectable or metastatic hepatocellular carcinoma; or with cisplatin and thiotepa to patients with primary or metastatic liver cancer.

10 In another embodiment, thalidomide is administered in combination with pegylated interferon alpha to patients with Kaposi's sarcoma.

In another embodiment, thalidomide is administered in combination with fludarabine, carboplatin, and/or topotecan to patients with refractory or relapsed or high-risk acute myelogenous leukemia.

15 In another embodiment, thalidomide is administered in combination with liposomal daunorubicin, topotecan and/or cytarabine to patients with unfavorable karyotype acute myeloblastic leukemia.

In another embodiment, thalidomide is administered in combination with gemcitabine and irinotecan to patients with non-small cell lung cancer. In one embodiment, thalidomide is administered in combination with carboplatin and irinotecan to patients with  
20 non-small cell lung cancer. In one embodiment, thalidomide is administered with doxorubicin to patients with non-small cell lung cancer who have been previously treated with carbo/VP  
16 and radiotherapy.

In another embodiment, thalidomide is administered in combination with carboplatin and/or taxotere, or in combination with carboplatin, paclitaxel and/or thoracic radiotherapy  
25 to patients with non-small cell lung cancer. In a specific embodiment, thalidomide is administered in combination with taxotere to patients with stage IIIB or IV non-small cell lung cancer.

In another embodiment, thalidomide is administered in combination with oblimersen (Genasense®) to patients with small cell lung cancer.

30 In another embodiment, thalidomide is administered alone or in combination with a second active ingredient such as vinblastine or fludarabine to patients with various types of lymphoma, including, but not limited to, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma or relapsed or refractory low grade follicular lymphoma.

In another embodiment, thalidomide is administered in combination with taxotere, IL-2, IFN, GM-CSF, and/or dacarbazine to patients with various types or stages of melanoma.

5 In another embodiment, thalidomide is administered alone or in combination with vinorelbine to patients with stage IIIB non-small cell lung cancer with pleural implants.

In another embodiment, thalidomide is administered to patients with various types or stages of multiple myeloma in combination with dexamethasone, zoledronic acid, palmitronate, GM-CSF, biaxin, vinblastine, melphalan, busulphan, cyclophosphamide, IFN, palmidronate, prednisone, bisphosphonate, celecoxib, arsenic trioxide, PEG INTRON-A,  
10 vincristine, or a combination thereof.

In another embodiment, thalidomide is administered to patients with relapsed or refractory multiple myeloma in combination with doxorubicin (Doxil<sup>®</sup>), vincristine and/or dexamethasone (Decadron<sup>®</sup>).

In another embodiment, thalidomide is administered to patients with various types or  
15 stages of ovarian cancer such as peritoneal carcinoma, papillary serous carcinoma, refractory ovarian cancer or recurrent ovarian cancer, in combination with taxol, carboplatin, doxorubicin, gemcitabine, cisplatin, xeloda, paclitaxel, dexamethasone, or a combination thereof.

In another embodiment, thalidomide is administered to patients with various types or  
20 stages of prostate cancer, in combination with xeloda, 5 FU/LV, gemcitabine, irinotecan plus gemcitabine, cyclophosphamide, vincristine, dexamethasone, GM-CSF, celecoxib, taxotere, ganciclovir, paclitaxel, adriamycin, docetaxel, estramustine, Emcyt, or a combination thereof.

In another embodiment, thalidomide is administered to patients with various types or  
25 stages of renal cell cancer, in combination with capecitabine, IFN, tamoxifen, IL-2, GM-CSF, Celebrex<sup>®</sup>, or a combination thereof.

In another embodiment, thalidomide is administered to patients with various types or stages of gynecologic, uterus or soft tissue sarcoma cancer in combination with IFN, a COX-2 inhibitor such as Celebrex<sup>®</sup>, and/or sulindac.

30 In another embodiment, thalidomide is administered to patients with various types or stages of solid tumors in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apocitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

In another embodiment, thalidomide is administered to patients with scleroderma or cutaneous vasculitis in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apocitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

5 This invention also encompasses a method of increasing the dosage of an anti-cancer drug or agent that can be safely and effectively administered to a patient, which comprises administering to a patient (*e.g.*, a human) thalidomide, or a pharmaceutically acceptable derivative, salt, solvate, clathrate, hydrate, or prodrug thereof. Patients that can benefit by this method are those likely to suffer from an adverse effect associated with anti-cancer drugs for treating a specific cancer of the skin, subcutaneous tissue, lymph nodes, brain,  
10 lung, liver, bone, intestine, colon, heart, pancreas, adrenal, kidney, prostate, breast, colorectal, or combinations thereof. The administration of thalidomide alleviates or reduces adverse effects which are of such severity that it would otherwise limit the amount of anti-cancer drug.

In one embodiment, thalidomide can be administered orally and daily in an amount  
15 of from about 50 to about 2,000 mg, preferably from about 50 to about 1,000 mg, and more preferably from about 50 to about 800 mg prior to, during, or after the occurrence of the adverse effect associated with the administration of an anti-cancer drug to a patient. In a particular embodiment, thalidomide is administered in combination with specific agents such as heparin, aspirin, coumadin, or G-CSF to avoid adverse effects that are associated  
20 with anti-cancer drugs such as but not limited to neutropenia or thrombocytopenia.

In one embodiment, thalidomide can be administered to patients with diseases and disorders associated with, or characterized by, undesired angiogenesis in combination with additional active ingredients including but not limited to anti-cancer drugs, anti-inflammatory, antihistamines, antibiotics, and steroids.

25 In another embodiment, this invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (*e.g.* before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based  
30 therapy presently used to treat, prevent or manage cancer. The combined use of thalidomide and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that thalidomide may provide additive or synergistic effects when given concurrently with conventional therapy.



In another embodiment, this invention encompasses a method of treating, preventing and/or managing diseases and disorders associated with, or characterized by, undesired angiogenesis, which comprises administering thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (e.g. before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based therapy presently used to treat, prevent or manage diseases and disorders associated with, or characterized by, undesired angiogenesis. The combined use of thalidomide and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that thalidomide may provide additive or synergistic effects when given concurrently with conventional therapy.

As discussed elsewhere herein, the invention encompasses a method of reducing, treating and/or preventing adverse or undesired effects associated with conventional therapy including, but not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy. Thalidomide and other active ingredient can be administered to a patient prior to, during, or after the occurrence of the adverse effect associated with conventional therapy.

In one embodiment, thalidomide can be administered in an amount of from about 50 to about 2,000 mg, preferably from about 50 to about 1,000 mg, and more preferably from about 50 to about 800 mg orally and daily alone, or in combination with a second active agent disclosed herein (*see, e.g.,* section 4.2), prior to, during, or after the use of conventional therapy.

In a specific embodiment of this method, thalidomide and doxorubicin are administered to patients with non-small cell lung cancer who were previously treated with carbo/VP 16 and radiotherapy.

#### 4.3.2 USE WITH TRANSPLANTATION THERAPY

Compounds of the invention can be used to reduce the risk of Graft Versus Host Disease (GVHD). Therefore, the invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with transplantation therapy.

As those of ordinary skill in the art are aware, the treatment of cancer is often based on the stages and mechanism of the disease. For example, as inevitable leukemic transformation develops in certain stages of cancer, transplantation of peripheral blood stem

cells, hematopoietic stem cell preparation or bone marrow may be necessary. The combined use of thalidomide and transplantation therapy provides a unique and unexpected synergism. In particular, thalidomide exhibits immunomodulatory activity that may provide additive or synergistic effects when given concurrently with transplantation therapy in  
5 patients with cancer.

Thalidomide can work in combination with transplantation therapy reducing complications associated with the invasive procedure of transplantation and risk of GVHD. This invention encompasses a method of treating, preventing and/or managing cancer which comprises administering to a patient (*e.g.*, a human) thalidomide, or a pharmaceutically  
10 acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow. Examples of stem cells suitable for use in the methods of the invention are disclosed in U.S. patent application no. 10/411,655, filed April 11, 2003 by R. Hariri *et al.*, the entirety of which is incorporated  
15 herein by reference.

In another embodiment, this invention encompasses a method of treating, preventing and/or managing diseases and disorders associated with, or characterized by, undesired angiogenesis, which comprises administering to a patient (*e.g.*, a human) thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug  
20 thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow.

In one embodiment of this method, thalidomide is administered to patients with multiple myeloma before, during, or after the transplantation of autologous peripheral blood progenitor cell.

25 In another embodiment, thalidomide is administered to patients with relapsing multiple myeloma after the stem cell transplantation.

In another embodiment, thalidomide and prednisone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous stem cell.

30 In another embodiment, thalidomide and dexamethasone are administered as salvage therapy for low risk post transplantation to patients with multiple myeloma.

In another embodiment, thalidomide and dexamethasone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous bone marrow.

In another embodiment, thalidomide is administered following the administration of high dose of melphalan and the transplantation of autologous stem cell to patients with chemotherapy responsive multiple myeloma.

5 In another embodiment, thalidomide and PEG INTRO-A are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous CD34-selected peripheral stem cell.

In another embodiment, thalidomide is administered with post transplant consolidation chemotherapy to patients with newly diagnosed multiple myeloma to evaluate anti-angiogenesis.

10 In another embodiment, thalidomide and dexamethasone are administered as maintenance therapy after DCEP consolidation, following the treatment with high dose of melphalan and the transplantation of peripheral blood stem cell to 65 years of age or older patients with multiple myeloma.

#### **4.3.3 CYCLING THERAPY.**

15 In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a patient. Cycling therapy involves the administration of an active agent for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or  
20 improves the efficacy of the treatment.

Consequently, in one specific embodiment of the invention, thalidomide is administered daily in a single or divided doses in a four to six week cycle with a rest period of about a week or two weeks. The invention further allows the frequency, number, and length of dosing cycles to be increased. Thus, another specific embodiment of the invention  
25 encompasses the administration of thalidomide for more cycles than are typical when it is administered alone. In yet another specific embodiment of the invention, thalidomide is administered for a greater number of cycles that would typically cause dose-limiting toxicity in a patient to whom a second active ingredient is not also being administered.

In one embodiment, thalidomide is administered daily and continuously for three or  
30 four weeks followed by a break of one or two weeks. In a particular embodiment, thalidomide is administered in an amount of from about 200 mg to about 800 mg/day for three to four weeks, followed by one week or two weeks of rest in a four or six week cycle.

In one embodiment of the invention, thalidomide and a second active ingredient are administered orally, with administration of thalidomide occurring 30 to 60 minutes prior to

a second active ingredient, during a cycle of four to six weeks. In another embodiment of the invention, the combination of thalidomide and a second active ingredient is administered by intravenous infusion over about 90 minutes every cycle. Typically, the number of cycles during which the combinatorial treatment is administered to a patient will  
5 be from about one to about 24 cycles, more typically from about two to about 16 cycles, and even more typically from about four to about three cycles.

#### 4.4 PHARMACEUTICAL COMPOSITIONS AND DOSAGE FORMS

Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms of the invention  
10 comprise thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. Pharmaceutical compositions and dosage forms of the invention can further comprise one or more excipients.

Pharmaceutical compositions and dosage forms of the invention can also comprise one or more additional active ingredients. Consequently, pharmaceutical compositions and  
15 dosage forms of the invention comprise the active ingredients disclosed herein (*e.g.*, thalidomide and a second active agent). Examples of optional second, or additional, active ingredients are disclosed herein (*see, e.g.*, section 4.2).

Single unit dosage forms of the invention are suitable for oral, mucosal (*e.g.*, nasal, sublingual, vaginal, buccal, or rectal), parenteral (*e.g.*, subcutaneous, intravenous, bolus  
20 injection, intramuscular, or intraarterial), topical (*e.g.*, eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (*e.g.*, nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal  
25 administration to a patient, including suspensions (*e.g.*, aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (*e.g.*, crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms  
30 suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the chronic treatment of the same disease. Similarly, a

parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which specific dosage forms encompassed by this invention will vary from one another will be readily apparent to those skilled in the art. See, e.g., *Remington's*  
5 *Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton PA (1990).

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form  
10 depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients may be  
15 accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or disaccharides. As used herein, the term "lactose-free" means that the amount of lactose  
20 present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the *U.S. Pharmacopeia* (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler,  
25 and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of  
30 some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, NY, 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds.

Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

5       Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

10       An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (*e.g.*, vials), blister packs,  
15       and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as "stabilizers," include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

20       Like the amounts and types of excipients, the amounts and specific types of active agents in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise thalidomide or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of from about 30 to about 2,000  
25       mg. Typical dosage forms comprise thalidomide or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of about 50, 100, 200, 300 or 400 mg. Typical dosage forms comprise the second active agent in an amount of from about 1 to about 3,500 mg, from about 5 to about 2,500 mg, from about 10 to about 500 mg, or from about 25 to about 250 mg. Of course, the specific amount of the second  
30       active agent will depend on the specific agent used, the type of disease being treated or managed, and the amount(s) of thalidomide and any optional additional active agents concurrently administered to the patient.

#### 4.4.1 ORAL DOSAGE FORMS

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (*e.g.*, chewable tablets), caplets, capsules, and liquids (*e.g.*, flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. *See generally, Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton PA (1990).

Typical oral dosage forms of the invention are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (*e.g.*, powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (*e.g.*, ethyl cellulose, cellulose acetate, carboxymethyl

cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (*e.g.*, Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. An specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103<sup>TM</sup> and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*, granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other algins, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (*e.g.*, peanut oil, cottonseed oil,



sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, TX), CAB-O-SIL  
5 (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, MA), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

A preferred solid oral dosage form of the invention comprises thalidomide,  
10 anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin. *See, e.g.,* U.S. Patent Application No. 10/608,077 filed June 30, 2003, the entirety of which is incorporated herein by reference.

#### **4.4.2 DELAYED RELEASE DOSAGE FORMS**

Active ingredients of the invention can be administered by controlled release means  
15 or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or  
20 more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the  
25 active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an  
30 optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other

characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

#### **4.4.3 PARENTERAL DOSAGE FORMS**

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention. For example, cyclodextrin and its derivatives can be used to increase the solubility of thalidomide and its derivatives. See, e.g., U.S. Patent No. 5,134,127, which is incorporated herein by reference.

#### 4.4.4 TOPICAL AND MUCOSAL DOSAGE FORMS

Topical and mucosal dosage forms of the invention include, but are not limited to, sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., *Remington's*  
5 *Pharmaceutical Sciences*, 16<sup>th</sup> and 18<sup>th</sup> eds., Mack Publishing, Easton PA (1980 & 1990); and *Introduction to Pharmaceutical Dosage Forms*, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used  
10 to provide topical and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and  
15 mixtures thereof to form solutions, emulsions or gels, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16<sup>th</sup> and 18<sup>th</sup> eds., Mack Publishing, Easton PA (1980 & 1990).

20 The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so  
25 as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-enhancing or penetration-enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

#### 4.4.5 KITS

30 Typically, active ingredients of the invention are preferably not administered to a patient at the same time or by the same route of administration. This invention therefore encompasses kits which, when used by the medical practitioner, can simplify the administration of appropriate amounts of active ingredients to a patient.

A typical kit of the invention comprises a dosage form of thalidomide, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, prodrug, or clathrate thereof. Kits encompassed by this invention can further comprise additional active ingredients such as oblimersen (Genasense<sup>®</sup>), melphalan, G-CSF, GM-CSF, EPO, 5 topotecan, dacarbazine, irinotecan, taxotere, IFN, COX-2 inhibitor, pentoxifylline, ciprofloxacin, dexamethasone, IL2, IL8, IL18, Ara-C, vinorelbine, isotretinoin, 13 cis-retinoic acid, or a pharmacologically active mutant or derivative thereof, or a combination thereof. Examples of the additional active ingredients include, but are not limited to, those disclosed herein (*see, e.g.*, section 4.2).

10 Kits of the invention can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

Kits of the invention can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active 15 ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous 20 vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl 25 benzoate.

## 5. EXAMPLE

One of the biological effects exerted by thalidomide is the reduction of synthesis of TNF- $\alpha$ . Thalidomide enhance the degradation of TNF- $\alpha$  mRNA. Inhibition of TNF- $\alpha$  production following LPS-stimulation of human PBMC by thalidomide was investigated *in* 30 *vitro*. The IC<sub>50</sub>'s of thalidomide for inhibiting production of TNF- $\alpha$  following LPS-stimulation of PBMC was ~194  $\mu$ M (50.1  $\mu$ g/mL).

The embodiments of the invention described above are intended to be merely exemplary, and those skilled in the art will recognize, or will be able to ascertain using no

more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures. All such equivalents are considered to be within the scope of the invention and are encompassed by the appended claims.